

Improved measurement of CP -violating parameters in $B^0 \rightarrow \rho^+ \rho^-$ decays

A. Somov,² A. J. Schwartz,² I. Adachi,⁷ H. Aihara,⁴² D. Anipko,¹ V. Aulchenko,¹ T. Aushev,^{16,11} A. M. Bakich,³⁷
 I. Bedny,¹ U. Bitenc,¹² I. Bizjak,¹² A. Bondar,¹ M. Bračko,^{7,18,12} T. E. Browder,⁶ M.-C. Chang,³ Y. Chao,²⁴
 A. Chen,²² K.-F. Chen,²⁴ W. T. Chen,²² B. G. Cheon,⁵ R. Chistov,¹¹ I.-S. Cho,⁴⁷ Y. Choi,³⁶ Y. K. Choi,³⁶
 S. Cole,³⁷ J. Dalseno,¹⁹ M. Dash,⁴⁶ J. Dragic,⁷ A. Drutskoy,² S. Eidelman,¹ S. Fratina,¹² G. Gokhroo,³⁸
 B. Golob,^{17,12} H. Ha,¹⁴ J. Haba,⁷ T. Hara,²⁹ K. Hayasaka,²⁰ H. Hayashii,²¹ M. Hazumi,⁷ D. Heffernan,²⁹
 T. Hokuue,²⁰ Y. Hoshi,⁴⁰ W.-S. Hou,²⁴ H. J. Hyun,¹⁵ T. Iijima,²⁰ K. Inami,²⁰ A. Ishikawa,⁴² H. Ishino,⁴³
 R. Itoh,⁷ M. Iwasaki,⁴² Y. Iwasaki,⁷ N. Joshi,³⁸ D. H. Kah,¹⁵ J. H. Kang,⁴⁷ N. Katayama,⁷ H. Kichimi,⁷
 H. O. Kim,³⁶ Y. J. Kim,⁴ K. Kinoshita,² S. Korpar,^{18,12} P. Križan,^{17,12} P. Krokovny,⁷ R. Kulasiri,²
 R. Kumar,³⁰ C. C. Kuo,²² A. Kuzmin,¹ Y.-J. Kwon,⁴⁷ J. S. Lee,³⁶ M. J. Lee,³⁵ S. E. Lee,³⁵ T. Lesiak,²⁵
 A. Limosani,⁷ S.-W. Lin,²⁴ D. Marlow,³² T. Matsumoto,⁴⁴ S. McOnie,³⁷ T. Medvedeva,¹¹ W. Mitaroff,⁹
 H. Miyake,²⁹ H. Miyata,²⁷ Y. Miyazaki,²⁰ R. Mizuk,¹¹ D. Mohapatra,⁴⁶ G. R. Moloney,¹⁹ M. Nakao,⁷ O. Nitoh,⁴⁵
 T. Nozaki,⁷ S. Ogawa,³⁹ T. Ohshima,²⁰ S. Okuno,¹³ S. L. Olsen,⁶ Y. Onuki,³³ H. Ozaki,⁷ P. Pakhlov,¹¹
 G. Pakhlova,¹¹ C. W. Park,³⁶ R. Pestotnik,¹² L. E. Piilonen,⁴⁶ H. Sahoo,⁶ Y. Sakai,⁷ O. Schneider,¹⁶ J. Schümann,⁷
 K. Senyo,²⁰ M. Shapkin,¹⁰ C. P. Shen,⁸ H. Shibuya,³⁹ B. Shwartz,¹ M. Starić,¹² H. Stoeck,³⁷ K. Sumisawa,⁷
 T. Sumiyoshi,⁴⁴ F. Takasaki,⁷ N. Tamura,²⁷ M. Tanaka,⁷ Y. Teramoto,²⁸ X. C. Tian,³¹ K. Trabelsi,⁷
 T. Tsukamoto,⁷ S. Uehara,⁷ K. Ueno,²⁴ T. Uglov,¹¹ Y. Unno,⁵ S. Uno,⁷ P. Urquijo,¹⁹ Y. Usov,¹ G. Varner,⁶
 K. E. Varvell,³⁷ S. Villa,¹⁶ A. Vinokurova,¹ C. C. Wang,²⁴ C. H. Wang,²³ M.-Z. Wang,²⁴ P. Wang,⁸
 Y. Watanabe,⁴³ E. Won,¹⁴ B. D. Yabsley,³⁷ A. Yamaguchi,⁴¹ Y. Yamashita,²⁶ Z. P. Zhang,³⁴ and A. Zupanc¹²

(The Belle Collaboration)

¹*Budker Institute of Nuclear Physics, Novosibirsk*

²*University of Cincinnati, Cincinnati, Ohio 45221*

³*Department of Physics, Fu Jen Catholic University, Taipei*

⁴*The Graduate University for Advanced Studies, Hayama*

⁵*Hanyang University, Seoul*

⁶*University of Hawaii, Honolulu, Hawaii 96822*

⁷*High Energy Accelerator Research Organization (KEK), Tsukuba*

⁸*Institute of High Energy Physics, Chinese Academy of Sciences, Beijing*

⁹*Institute of High Energy Physics, Vienna*

¹⁰*Institute of High Energy Physics, Protvino*

¹¹*Institute for Theoretical and Experimental Physics, Moscow*

¹²*J. Stefan Institute, Ljubljana*

¹³*Kanagawa University, Yokohama*

¹⁴*Korea University, Seoul*

¹⁵*Kyungpook National University, Taegu*

¹⁶*Swiss Federal Institute of Technology of Lausanne, EPFL, Lausanne*

¹⁷*University of Ljubljana, Ljubljana*

¹⁸*University of Maribor, Maribor*

¹⁹*University of Melbourne, School of Physics, Victoria 3010*

²⁰*Nagoya University, Nagoya*

²¹*Nara Women's University, Nara*

²²*National Central University, Chung-li*

²³*National United University, Miao Li*

²⁴*Department of Physics, National Taiwan University, Taipei*

²⁵*H. Niewodniczanski Institute of Nuclear Physics, Krakow*

²⁶*Nippon Dental University, Niigata*

²⁷*Niigata University, Niigata*

²⁸*Osaka City University, Osaka*

²⁹*Osaka University, Osaka*

³⁰*Panjab University, Chandigarh*³¹*Peking University, Beijing*³²*Princeton University, Princeton, New Jersey 08544*³³*RIKEN BNL Research Center, Upton, New York 11973*³⁴*University of Science and Technology of China, Hefei*³⁵*Seoul National University, Seoul*³⁶*Sungkyunkwan University, Suwon*³⁷*University of Sydney, Sydney, New South Wales*³⁸*Tata Institute of Fundamental Research, Mumbai*³⁹*Toho University, Funabashi*⁴⁰*Tohoku Gakuin University, Tagajo*⁴¹*Tohoku University, Sendai*⁴²*Department of Physics, University of Tokyo, Tokyo*⁴³*Tokyo Institute of Technology, Tokyo*⁴⁴*Tokyo Metropolitan University, Tokyo*⁴⁵*Tokyo University of Agriculture and Technology, Tokyo*⁴⁶*Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061*⁴⁷*Yonsei University, Seoul*

We present a measurement of the CP -violating asymmetry in $B^0 \rightarrow \rho^+ \rho^-$ decays using 535 million $B\bar{B}$ pairs collected with the Belle detector at the KEKB e^+e^- collider. We measure CP -violating coefficients $\mathcal{A} = 0.16 \pm 0.21$ (stat) ± 0.08 (syst) and $\mathcal{S} = 0.19 \pm 0.30$ (stat) ± 0.08 (syst). These values are used to determine the unitarity triangle angle ϕ_2 using an isospin analysis; the solution consistent with the Standard Model lies in the range $54^\circ < \phi_2 < 113^\circ$ at the 90% confidence level.

PACS numbers: 13.25.Hw, 12.15.Hh, 11.30.Er

CP violation in the Standard Model is attributed to the presence of an irreducible complex phase in the Cabibbo-Kobayashi-Maskawa [1] (CKM) quark-mixing matrix. The unitarity of the CKM matrix leads to six triangles in the complex plane. One such triangle is given by the following relation among the matrix elements: $V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* = 0$. The phase angle ϕ_2 , defined as $\arg[-(V_{td}V_{tb}^*)/(V_{ud}V_{ub}^*)]$, can be determined by measuring a time-dependent CP asymmetry in $b \rightarrow u\bar{d}$ decays such as $B^0 \rightarrow \pi^+\pi^-$, $(\rho\pi)^0$, and $\rho^+\rho^-$ [2]. The time-dependent rate for $B \rightarrow \rho^+\rho^-$ decays tagged with $B^0(q = +1)$ and $\bar{B}^0(q = -1)$ mesons is given by

$$\mathcal{P}_{\rho\rho}(\Delta t) = \frac{e^{-|\Delta t|/\tau_{B^0}}}{4\tau_{B^0}} \{1 + q[\mathcal{A} \cos(\Delta m \Delta t) + \mathcal{S} \sin(\Delta m \Delta t)]\}, \quad (1)$$

where τ_{B^0} is the B^0 lifetime, Δm is the mass difference between the two B^0 mass eigenstates, Δt is the proper-time difference between the two B decays in the event, and $\mathcal{A}$ and $\mathcal{S}$ are CP asymmetry coefficients. If the decay amplitude is a pure CP -even state and is dominated by a tree diagram, $\mathcal{S} = \sin(2\phi_2)$ and $\mathcal{A} = 0$. The presence of an amplitude with a different weak phase (such as from a loop diagram) gives rise to direct CP violation and shifts $\mathcal{S}$ from $\sin(2\phi_2)$. However, the size of a loop amplitude is constrained to be small by the small branching fraction of $B^0 \rightarrow \rho^0 \rho^0$ [3].

The CP -violating parameters receive contributions from a longitudinally polarized state (CP -even) and two transversely polarized states (an admixture of CP -even and CP -odd states). Recent measurements of the polarization fraction by Belle [4] and BaBar [5] show that

the longitudinal polarization fraction is near unity ($f_L = 0.968 \pm 0.023$ [6]).

Here we present an improved measurement of the CP -violating coefficients $\mathcal{A}$ and $\mathcal{S}$ using 492 fb $^{-1}$ of data containing 535 million $B\bar{B}$ pairs. This data sample is about a factor of two larger than that used in our earlier publication [4]. In addition we have modified the event selection by reducing the threshold on a continuum suppression variable; this increases our reconstruction efficiency by about 70%. We subsequently introduce a probability density function (PDF) for the continuum suppression variable into the likelihood function; this provides additional discrimination between signal and backgrounds. The expected improvement in the statistical error of $\mathcal{A}$ and $\mathcal{S}$ due to the new event selection is about 12%.

The $B\bar{B}$ pairs were collected with the Belle detector [7] at the KEKB [8] e^+e^- asymmetric-energy (3.5 GeV on 8.0 GeV) collider with a center-of-mass (CM) energy at the $\Upsilon(4S)$ resonance. The $\Upsilon(4S)$ is produced with a Lorentz boost of $\beta\gamma = 0.425$ nearly along the z axis, which is oriented antiparallel to the positron beam. Since the B^0 and $\bar{B}^0$ mesons are produced approximately at rest in the $\Upsilon(4S)$ CM system, the decay time difference Δt is related to the distance between the decay vertices of the two B mesons as $\Delta t \simeq \Delta z/\beta\gamma c$, where c is the speed of light.

The Belle detector [7] is a large-solid-angle spectrometer. It includes a silicon vertex detector (SVD), a 50-layer central drift chamber (CDC), an array of aerogel threshold Cherenkov counters (ACC), time-of-flight scintillation counters (TOF), and an electromagnetic calorimeter (ECL) comprised of CsI(Tl) crystals located inside a




FIG. 1: Left: projections in M_{bc} for events satisfying $-0.10 \text{ GeV} < \Delta E < 0.06 \text{ GeV}$. Right: projections in ΔE for events with $5.27 \text{ GeV}/c^2 < M_{bc} < 5.29 \text{ GeV}/c^2$. The top plots correspond to good quality tags ($0.75 < r < 1.0$), and the bottom plots correspond to lower quality tags ($r < 0.75$). The curves show fit projections: dashed is $\rho^+\rho^- + \rho\pi\pi$, dotted is $q\bar{q}$, dot-dashed is $b \rightarrow c$, small solid is $b \rightarrow u$, and large solid is the total. For these plots the $\mathcal{R}$ requirement has been tightened to increase the ratio of signal to background.

superconducting solenoid coil that provides a 1.5 T magnetic field.

We reconstruct $B^0 \rightarrow \rho^+\rho^-$ decays by combining two oppositely charged pion tracks with two neutral pions. Each charged track is required to have a transverse momentum $p_T > 0.10 \text{ GeV}/c$ in the laboratory frame and originate within $dr < 0.2 \text{ cm}$ in the radial direction and within $|dz| < 4.0 \text{ cm}$ along the z -axis from the interaction point, which is determined run-by-run. A track is identified as a pion using information from the CDC, ACC and TOF systems. Tracks matched with clusters in the ECL that are consistent with an electron hypothesis are rejected.

The π^0 candidates are reconstructed from $\gamma\gamma$ pairs with an invariant mass in the range $117.8 \text{ MeV}/c^2 < M_{\gamma\gamma} < 150.2 \text{ MeV}/c^2$ (about $\pm 3\sigma$ in m_{π^0} resolution). Photons are required to have energy $E_\gamma > 50 \text{ MeV}$ in the ECL barrel region ($32^\circ < \theta < 129^\circ$) and $E_\gamma > 90 \text{ MeV}$ in the endcap regions ($17^\circ < \theta < 32^\circ$ and $129^\circ < \theta < 150^\circ$), where θ denotes the polar angle with respect to the z axis.

To reconstruct $\rho^\pm$ mesons, we combine $\pi^\pm$ candidates with π^0 candidates. The $\pi^\pm\pi^0$ combination must have an invariant mass in the range $0.62 \text{ GeV}/c^2 < M_{\pi^\pm\pi^0} < 0.92 \text{ GeV}/c^2$. To reduce combinatorial background, we reject π^0 's with $p < 0.35 \text{ GeV}/c$ in the CM frame. We




FIG. 2: $\mathcal{R}$ distribution for high-purity tagged events satisfying $5.27 \text{ GeV}/c^2 < M_{bc} < 5.29 \text{ GeV}/c^2$ and $-0.10 \text{ GeV} < \Delta E < 0.06 \text{ GeV}$. The curves show fit projections: dashed is $\rho^+\rho^- + \rho\pi\pi$, dotted is $q\bar{q}$, dot-dashed is $b \rightarrow c$, small solid is $b \rightarrow u$, and large solid is the total.

also require $-0.80 < \cos \theta_\pm < 0.98$, where $\theta_\pm$ is the angle between the direction of the π^0 from the $\rho^\pm$ and the negative of the B^0 momentum in the $\rho^\pm$ rest frame.

$B^0 \rightarrow \rho^+\rho^-$ decays are identified using the beam-energy-constrained mass $M_{bc} \equiv \sqrt{E_{\text{beam}}^2 - p_B^2}$ and energy difference $\Delta E \equiv E_B - E_{\text{beam}}$, where E_{beam} is the beam energy, and E_B and p_B are the energy and momentum of the reconstructed B candidate, all evaluated in the CM frame.

The flavor of the B meson accompanying the $B^0 \rightarrow \rho^+\rho^-$ candidate is identified via its decay products: charged leptons, kaons, and Λ 's. A tagging algorithm [9] yields the flavor of the tagged meson, q , and a quality factor, r . The parameter r ranges from 0 for no flavor discrimination to 1 for unambiguous flavor assignment. We divide the data sample into six r intervals (denoted $\ell=1,2,\dots,6$). The wrong tag fractions ω_ℓ for these intervals and the differences $\Delta\omega_\ell$ in these fractions between B^0 and $\bar{B}^0$ decays are determined from data [9].

The dominant background originates from $e^+e^- \rightarrow q\bar{q}$ ($q = u, d, s, c$) continuum events. To separate $q\bar{q}$ jet-like events from more spherical $B\bar{B}$ events, we use event-shape variables, specifically, 16 modified Fox-Wolfman moments [11] combined into a Fisher discriminant [12]. We form signal and background likelihood functions $\mathcal{L}_s$ and $\mathcal{L}_{BG}$ by multiplying the PDF for the Fisher discriminant by a PDF for $\cos \theta_B$, where θ_B is the polar angle in the CM frame between the B direction and the beam axis. The PDFs for signal and $q\bar{q}$ are obtained from Monte Carlo (MC) simulations and the data sideband $5.23 \text{ GeV}/c^2 < M_{bc} < 5.26 \text{ GeV}/c^2$, respectively. We calculate the ratio $\mathcal{R} = \mathcal{L}_s/(\mathcal{L}_s + \mathcal{L}_{BG})$ and make a loose requirement $\mathcal{R} > 0.15$.

The decay vertices of a $\rho^+\rho^-$ candidate and the tag-side B meson are reconstructed using charged tracks that have a sufficient number of SVD hits and an interaction point constraint. The vertex reconstruction algorithm is described in Ref. [10].

The analysis is organized in two main steps. We first

determine the yields of signal and background components from a fit to the three-dimensional $(M_{bc}, \Delta E, \mathcal{R})$ distribution. Here, B^0 candidates are required to satisfy $5.23 \text{ GeV}/c^2 < M_{bc} < 5.29 \text{ GeV}/c^2$, $-0.2 \text{ GeV} < \Delta E < 0.26 \text{ GeV}$, and $\mathcal{R} > 0.15$. We subsequently perform a fit to the Δt distribution to determine the CP parameters $\mathcal{A}$ and $\mathcal{S}$. The signal region used for the Δt fit is $5.27 \text{ GeV}/c^2 < M_{bc} < 5.29 \text{ GeV}/c^2$, $-0.12 \text{ GeV} < \Delta E < 0.08 \text{ GeV}$, and $\mathcal{R} > 0.15$.

About 12.6% of events contain multiple $B^0 \rightarrow \rho^+ \rho^-$ candidates, most of which arise from fake π^0 's combining with good tracks. We select the best candidate based on the π^0 masses, i.e., minimizing $\sum_{\pi_{1,2}^0} (m_{\gamma\gamma} - m_{\pi^0})^2$. For the small fraction (3%) of multiple-candidate events that arise due to extraneous $\pi^\pm$ tracks combining with a single π^0 , we select one randomly. Signal decays that have at least one π meson incorrectly identified are referred to as ‘‘self-cross-feed’’ (SCF) events.

The likelihood function used to determine the event yields is given by

$$\mathcal{L} = \exp \left(- \sum_j N_j \right) \prod_{i=1}^{N_{\text{evt}}} \left[\sum_j N_j \mathcal{P}_j(M_{bc}^i, \Delta E^i, \mathcal{R}^i) \right], \quad (2)$$

where j indicates one of the following event categories: signal and $\rho\pi\pi$ non-resonant decays, SCF events, continuum background ($q\bar{q}$), $b \rightarrow c$ background, and charmless ($b \rightarrow u$) background. N_j is the yield of each category, $\mathcal{P}_j(M_{bc}^i, \Delta E^i, \mathcal{R}^i)$ is the PDF for the i -th event for category j , and N_{evt} is the total number of events in the fit. Except for the small contributions of $b \rightarrow u$ background and SCF events, the yields N_j are determined from the fit. Due to the similar shapes of the M_{bc} , ΔE , and $\mathcal{R}$ distributions for signal and $\rho\pi\pi$ events, we cannot distinguish these two components; the fraction of $\rho\pi\pi$ events is measured in Ref. [4] and constitutes $(6.3 \pm 6.7)\%$ of the total $N_{\rho^+ \rho^- + \rho\pi\pi}$ signal. The fraction of SCF events is determined from MC simulation. The M_{bc} and ΔE shapes for the signal and SCF components are modeled by a two-dimensional smoothed histogram obtained from a large MC sample. To take into account a small difference between the MC and data, the M_{bc} - ΔE shapes are corrected according to calibration factors determined from a $B^+ \rightarrow \bar{D}^0 \rho^+$, $\bar{D}^0 \rightarrow K^+ \pi^- \pi^0$ control sample. The $\mathcal{R}$ shapes are modeled by one-dimensional histograms, also obtained from MC simulation.

The PDF for $b \rightarrow c$ background is the product of a threshold ARGUS function [13] for M_{bc} , a quadratic polynomial for ΔE , and the sum of a Gaussian and a third-order polynomial for $\mathcal{R}$. The shapes of the ΔE and $\mathcal{R}$ distributions depend on the tag quality bin ℓ . Parameters for all distributions are obtained from the MC.

The M_{bc} and ΔE PDFs for $q\bar{q}$ are modeled by an ARGUS function and a linear function, respectively. The ΔE slope depends on $\mathcal{R}$ and the tag quality bin ℓ . The

shape parameters for M_{bc} and ΔE are floated in the fit. The $\mathcal{R}$ PDF for $q\bar{q}$ background is taken to be an eighth-order polynomial function; the coefficients depend on the bin ℓ and are determined from a data sample collected at a CM energy $\sim 60 \text{ MeV}$ below the $\Upsilon(4S)$.

The $b \rightarrow u$ background is dominated by $B \rightarrow (\rho\pi, a_1\pi, a_1\rho)$ decays. We estimate the $B^\pm \rightarrow (a_1\pi)^\pm$ branching fractions (which are unmeasured) to be $(20 \pm 10) \times 10^{-6}$ using the measured value for $B^0 \rightarrow a_1^\pm \pi^\mp$ [14]. For $B^\pm \rightarrow (a_1\rho)^\pm$ we assume branching fractions of $(30 \pm 15) \times 10^{-6}$, consistent with the present upper limit for $B^0 \rightarrow a_1^\pm \rho^\mp$ ($< 6 \times 10^{-5}$ [15]). The fraction of $b \rightarrow u$ events is very small (0.37%) and thus is fixed in the fit according to the prediction of MC simulation. A fit to 176843 events maximizing $\mathcal{L}$ yields $N_{\rho\rho+\rho\pi\pi} = 576 \pm 53$. Figures 1 and 2 show the M_{bc} , ΔE , and $\mathcal{R}$ distributions along with projections of the fit result.

The CP -violating parameters $\mathcal{A}$ and $\mathcal{S}$ are obtained using an unbinned ML fit to the Δt distribution. The likelihood function for event i is given by

$$\mathcal{L}_i = \sum_n \int f_n(\vec{x}_i) \mathcal{P}_n(\Delta t') R_n(\Delta t^i, \Delta t') d\Delta t', \quad (3)$$

where n is one of the six event categories: correctly reconstructed signal, SCF events, $\rho\pi\pi$ non-resonant events, $b \rightarrow c$ background, $q\bar{q}$ background, and $b \rightarrow u$ background. The weights f_n are functions of $\vec{x} \in (M_{bc}, \Delta E, \mathcal{R})$ and are normalized to the event fractions obtained from the $(M_{bc}, \Delta E, \mathcal{R})$ fit. The PDFs $\mathcal{P}_n(\Delta t)$ are convolved with the corresponding Δt resolution functions R_n . Both f_n and $\mathcal{P}_n(\Delta t)$ depend on the tag quality bin ℓ .

The signal PDF is given by Eq. (1) modified to take into account the effect of incorrect flavor assignment: $e^{-|\Delta t|/\tau_{B^0}}/(4\tau_{B^0}) \times \{1 - q\Delta\omega_\ell + q(1 - 2\omega_\ell) [\mathcal{A} \cos(\Delta m \Delta t) + \mathcal{S} \sin(\Delta m \Delta t)]\}$. As the fraction of longitudinal polarization f_L is close to 100%, we assume that $\mathcal{A} = \mathcal{A}_L$, $\mathcal{S} = \mathcal{S}_L$, and consider the potential contribution from a transversely polarized amplitude as a systematic uncertainty. The signal PDF is convolved with the same Δt resolution function as that used for Belle's $\sin 2\phi_1$ measurement [10].

The fraction of SCF events with incorrectly reconstructed vertices is estimated from MC simulation to be $(6.5 \pm 0.1)\%$ of all signal events. The PDFs $\mathcal{P}_{\rho\pi\pi}$ and $\mathcal{P}_{\text{SCF}}$ are exponential with $\tau = \tau_B$ and $\tau \approx 0.96 \text{ ps}$ (from MC), respectively; these are smeared by a common resolution function.

The Δt PDFs for the backgrounds are modeled as a sum of prompt and exponential components: $\mathcal{P}_k = f_\delta^k \delta(\Delta t) + (1 - f_\delta^k) e^{-|\Delta t|/\tau_k}/2\tau_k$, where k represents continuum, $b \rightarrow c$, and $b \rightarrow u$ backgrounds, f_δ^k is the fraction of the prompt component, $\delta(\Delta t)$ is the Dirac delta function, and τ_k is an effective lifetime. These PDFs are convolved with a resolution-like function R_k parameterized




FIG. 3: The Δt distribution and projections of the fit for events satisfying $0.5 < r < 1.0$: (a) $q = +1$ tags, (b) $q = -1$ tags. The hatched region shows signal events. The raw CP asymmetry is shown in (c). For these plots the $\mathcal{R}$ requirement has been tightened to increase the ratio of signal to background.

as a sum of two Gaussian functions. Parameters for $\mathcal{P}_k$ and R_k are determined from a data sideband for continuum background and from large MC samples for $b \rightarrow c$ and $b \rightarrow u$ backgrounds. To account for small correlations between the shape of the Δt distribution and $\mathcal{R}$ for $q\bar{q}$ background, the parameters are obtained separately for low ($0.15 < \mathcal{R} < 0.75$) and high ($0.75 < \mathcal{R} < 1.0$) $\mathcal{R}$ regions.

We determine $\mathcal{A}$ and $\mathcal{S}$ by maximizing $\sum_i \log \mathcal{L}_i$, where i runs over the 18016 events in the $(M_{bc}, \Delta E, \mathcal{R})$ signal region. The results are $\mathcal{A} = 0.16 \pm 0.21$ and $\mathcal{S} = 0.19 \pm 0.30$, where the errors are statistical. The correlation coefficient is -0.10 . These values are consistent with no CP violation ($\mathcal{A} = \mathcal{S} = 0$); the errors are consistent with MC expectations. Figure 3 shows the data and projections of the fit result.

The sources of systematic error are listed in Table I. The error for most sources is evaluated by varying the corresponding parameters by ± 1 standard deviation (σ). The effect of a possible asymmetry in $b \rightarrow c$ and $q\bar{q}$ is evaluated by adding such an asymmetry to the $b \rightarrow c$ and $q\bar{q}$ Δt distributions. The uncertainty due to a possible asymmetry in $\rho\pi\pi$ non-resonant decays is estimated by varying $\mathcal{A}_{\rho\pi\pi}$ and $\mathcal{S}_{\rho\pi\pi}$ by 0.68, corresponding to a 68% confidence interval of a free distribution. We vary the branching fractions for $a_1\rho$ and $a_1\pi$ decays and also allow for a CP asymmetry of up to 100% in these modes. The




FIG. 4: $1 - \text{C.L.}$ vs. ϕ_2 . The horizontal lines denote C.L. = 68.3% (solid) and C.L. = 90% (dashed).

error due to transverse polarization is obtained by first setting f_L equal to its central value [6] and varying $\mathcal{A}_T$, $\mathcal{S}_T$ from -1 to $+1$; then, conservatively assuming that the transversely polarized component (with fraction $f_T = 1 - f_L$) is pure CP -odd for which $\mathcal{A}_T = \mathcal{A}_L$, $\mathcal{S}_T = -\mathcal{S}_L$, and varying f_L by its error. Summing up in quadrature all systematic uncertainties, we obtain overall systematic errors of ± 0.08 for both $\mathcal{A}$ and $\mathcal{S}$. Thus,

$$\mathcal{A} = 0.16 \pm 0.21 (\text{stat}) \pm 0.08 (\text{syst}) \quad (4)$$

$$\mathcal{S} = 0.19 \pm 0.30 (\text{stat}) \pm 0.08 (\text{syst}). \quad (5)$$

These values are consistent with, and supersede, our previous measurement [4]. They are also consistent with results obtained by BaBar [5].

We constrain ϕ_2 using an isospin analysis [17], which allows one to relate six observables to six underlying parameters: five decay amplitudes for $B \rightarrow \rho\rho$ and the angle ϕ_2 . The observables are the branching fractions for $B \rightarrow \rho^+\rho^-$, $\rho^+\rho^0$ [6], and $\rho^0\rho^0$ [3]; the CP parameters $\mathcal{A}$ and $\mathcal{S}$ (our results); and the parameter $\mathcal{A}_{\rho^0\rho^0}$ for $B \rightarrow \rho^0\rho^0$ decays. The last parameter is not yet measured, but nevertheless one can constrain ϕ_2 . The branching fractions must be multiplied by the corresponding longitudinal polarization fractions [6]. We neglect possible contributions from electroweak penguins and $I = 1$ amplitudes [18] and possible interference between signal and non-resonant components. We follow the statistical method of Ref. [19] and construct a $\chi^2(\phi_2)$ using the measured values and obtain a minimum χ^2 (denoted $\chi^2_{\min}$); we then scan ϕ_2 from 0° to 180° , calculating the difference $\Delta\chi^2 \equiv \chi^2(\phi_2) - \chi^2_{\min}$. We insert $\Delta\chi^2$ into the cumulative distribution function for the χ^2 distribution for one degree of freedom to obtain a confidence level (C.L.) for each ϕ_2 value. The resulting function $1 - \text{C.L.}$ (Fig. 4) has more than one peak due to ambiguities that arise when solving for ϕ_2 . The “flat-top” regions in Fig. 4

TABLE I: Systematic errors for CP coefficients $\mathcal{A}$ and $\mathcal{S}$.

Type	$\Delta\mathcal{A} (\times 10^{-2})$		$\Delta\mathcal{S} (\times 10^{-2})$	
	$+\sigma$	$-\sigma$	$+\sigma$	$-\sigma$
Wrong tag fractions	0.5	0.5	0.8	0.8
Parameters $\Delta m, \tau_{B^0}$	0.2	0.3	0.6	0.7
Resolution function	1.4	1.5	1.0	1.7
Background Δt distributions	0.5	0.5	1.0	1.1
Component fractions	1.5	1.9	3.9	3.7
$\rho\pi\pi$ non-resonant fractions	1.2	1.0	1.5	1.2
SCF fraction, Δt PDF	0.2	0.2	0.1	0.1
Shape of $\mathcal{R}$ PDF	0.8	0.7	1.2	1.3
Vertexing	2.1	2.1	1.0	1.3
Possible fitting bias	0.3	0.0	0.3	0.0
Background asymmetry	1.1	0.0	0.0	0.4
$b \rightarrow u$ asymmetry	2.4	2.9	2.4	3.2
$\rho\pi\pi$ asymmetry	4.6	4.6	4.6	4.6
Transverse polarization	3.8	2.8	4.6	2.7
Tag-side interference [16]	3.7	3.7	0.1	0.1
Total	8.3	8.0	8.4	7.9

arise because $\mathcal{A}_{\rho^0\rho^0}$ is not measured. The solution consistent with the Standard Model is $62^\circ < \phi_2 < 106^\circ$ at 68% C.L. or $54^\circ < \phi_2 < 113^\circ$ at 90% C.L. Recently, an alternative model-dependent approach to extract ϕ_2 using flavor $SU(3)$ symmetry has been proposed [20]. This method could potentially give more stringent constraints on ϕ_2 .

In summary, we present an improved measurement of the CP -violating coefficients $\mathcal{A}$ and $\mathcal{S}$ in $B^0 \rightarrow \rho^+ \rho^-$ decays using 492 fb^{-1} of data, which corresponds to 535 million $B\bar{B}$ pairs. These measurements are used to constrain the angle ϕ_2 .

We thank the KEKB group for the excellent operation of the accelerator, the KEK cryogenics group for the efficient operation of the solenoid, and the KEK computer group and the National Institute of Informatics for valuable computing and Super-SINET network support. We acknowledge support from the Ministry of Education, Culture, Sports, Science, and Technology of Japan and the Japan Society for the Promotion of Science; the Australian Research Council and the Australian Department of Education, Science and Training; the National Science Foundation of China and the Knowledge Innovation Program of the Chinese Academy of Sciences under contract No. 10575109 and IHEP-U-503; the Department of Science and Technology of India; the BK21 program of the

Ministry of Education of Korea, the CHEP SRC program and Basic Research program (grant No. R01-2005-000-10089-0) of the Korea Science and Engineering Foundation, and the Pure Basic Research Group program of the Korea Research Foundation; the Polish State Committee for Scientific Research; the Ministry of Education and Science of the Russian Federation and the Russian Federal Agency for Atomic Energy; the Slovenian Research Agency; the Swiss National Science Foundation; the National Science Council and the Ministry of Education of Taiwan; and the U.S. Department of Energy.

-
- [1] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973); N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963).
 - [2] Charge-conjugate modes are included throughout this paper unless noted otherwise.
 - [3] B. Aubert *et al.* (BABAR Collaboration), hep-ex/0612021 (2006), submitted to Phys. Rev. Lett.
 - [4] A. Somov *et al.* (Belle Collaboration), Phys. Rev. Lett. **96**, 171801 (2006).
 - [5] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. Lett. **95**, 041805 (2005); Phys. Rev. Lett. **93**, 231801 (2004); hep-ex/0607098.
 - [6] Heavy Flavor Averaging Group, August 2006, <http://www.slac.stanford.edu/xorg/hfag/>.
 - [7] A. Abashian *et al.* (Belle Collaboration), Nucl. Instr. Meth. A **479**, 117 (2002).
 - [8] S. Kurokawa and E. Kikutani, Nucl. Instr. and Meth. A **499**, 1 (2003), and other papers in this volume.
 - [9] H. Kakuno *et al.*, Nucl. Instr. Meth. A **533**, 516 (2004).
 - [10] K.-F. Chen *et al.* (Belle Collaboration), Phys. Rev. D **72**, 012004 (2005); H. Tajima *et al.*, Nucl. Instr. Meth. A **533** (2004).
 - [11] G. C. Fox and S. Wolfram, Phys. Rev. Lett. **41**, 1581 (1978).
 - [12] S. H. Lee *et al.*, Phys. Rev. Lett. **91**, 261801 (2003).
 - [13] H. Albrecht *et al.* (Argus Collaboration), Phys. Lett. B **241**, 278 (1990).
 - [14] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. Lett. **97**, 051802 (2006); K. Abe *et al.* (Belle Collaboration), hep-ex/0507096.
 - [15] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **74**, 031104 (2006).
 - [16] O. Long *et al.*, Phys. Rev. D **68**, 034010 (2003).
 - [17] M. Gronau and D. London, Phys. Rev. Lett. **65**, 3381 (1990).
 - [18] A. Falk *et al.*, Phys. Rev. D **69**, 011502(R) (2004).
 - [19] J. Charles *et al.* (CKMfitter Group), Eur. Phys. J. C **41**, 1 (2005).
 - [20] M. Beneke, M. Gronau, J. Rohrer and M. Spranger, Phys. Lett. B **638**, 68 (2006).